arXiv:1401.4110v1 [physics.atom-ph] 16 Jan 2014

Radiative rates for E1, E2, M1, and M2 transitions among the $3s^2 3p^5$, $3s3p^6$, and $3s^2 3p^4 3d$ configurations of Cl-like W LVIII

Kanti M Aggarwal and Francis P Keenan

Astrophysics Research Centre, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, Northern Ireland, UK

e-mail: K.Aggarwal@qub.ac.uk

Received 13 November 2013. Accepted 16 January 2014

PACS Ref: 31.25 Jf, 32.70 Cs, 95.30 Ky

Abstract

We report calculations of energy levels, radiative decay rates, and lifetimes for transitions among the $3s^2 3p^5$, $3s3p^6$, and $3s^2 3p^4 3d$ configurations of Cl-like W LVIII. The general-purpose relativistic atomic structure package (GRASP) has been adopted for our calculations. Comparisons are made with the most recent results of Mohan et al. [Can. J. Phys. **92** (2014) xxx] and discrepancies in lifetimes are noted, up to four orders of magnitude in some instances. Our energy levels are estimated to be accurate to better than 0.5%, whereas results for radiative rates and lifetimes should be accurate to better than 20%.

1 Introduction

Atomic data, including energy levels and oscillator strengths (or radiative decay rates), for tungsten (W) ions have been of great interest, both theoretically [1]–[2] and experimentally [3]–[5], for several years. This is mainly because tungsten is one of the constituents of tokamak reactor walls. Due to high temperatures of such fusion plasmas, many ionisation stages of W are observable and atomic data are required to assess radiation losses. The need for atomic data for W ions has become more urgent due to the developing ITER project. However, there are two requirements from the user community, namely the data should be generated for a significantly large model [6] and be reliable [7]. A complete set of data from a large model takes account of cascading effects, while reliability (based on rigorous tests) provides confidence to the users.

In a recent paper, Mohan et al. [8] have reported results for energy levels, oscillator strengths, radiative rates, line strengths, and lifetimes for Cl-like W LVIII. For the calculations, they adopted the widely used and readily available GRASP (general-purpose relativistic atomic structure package) code. This was originally developed by Grant et al. [9] and has undergone several revisions by the names GRASP [10], GRASP2 [11], and GRASP2K [12]–[13]. The version they used has been revised by one of its authors (Dr. P. H. Norrington), is known as GRASP0, and is available at `http://web.am.qub.ac.uk/DARC/`. It is a fully relativistic code, based on the jj coupling scheme. Further relativistic corrections arising from the Breit interaction and QED (quantum electrodynamics) effects have also been included. Furthermore, this version provides compatible results with others.

Tungsten is a heavy element (Z = 74) and therefore relativistic effects are very important in the determination of its atomic structure. However, equally important is the inclusion of *configuration interaction* (CI) – see for example, Fournier [1] and Aggarwal and Keenan [14]. In particular, Cl-like ions are very complicated and CI is highly important, as demonstrated by Aggarwal and Keenan for Ti VI [15], Cr VIII [16], Fe X [17], and Co XI [18]. These are comparatively light ions but W LVIII is no exception, and therefore Mohan et al. [8] have included CI among 15 configurations (GRASP1), namely $3s^23p^5$, $3s3p^6$, $3s^23p^43d$, $3s^23p^44\ell$, $3s3p^53d$, $3s^23p^33d^2$, $3s3p^43d^2$, $3p^63d$, $3s^23p^23d^3$, $3p^53d^2$, $3s^23p^34d^2$, and $3s^23p^34f^2$. However, inclusion of this limited CI is not sufficient for an accurate determination of energy levels and radiative rates (A- values). Furthermore, they reported energies for only 31 levels of the $3s^23p^5$, $3s3p^6$, and $3s^23p^43d$ configurations, and A- values for only transitions from the two levels of the ground state, i.e. $(3s^23p^5)\ {}^2P^o_{3/2,1/2}$. Therefore, there is scope for improvement and to extend the range of transitions.

2 Energy levels

For our calculations we have employed the same GRASP code as adopted by Mohan et al. [8]. Additionally, we have used the option of *extended average level* (EAL), in which a weighted (proportional to $2j+1$) trace of the Hamiltonian matrix is minimised. This produces a compromise set of orbitals comparable to other options, such as *average level* (AL), as noted by Aggarwal et al. for several ions of Kr [19] and Xe [20]. However, in our calculations we include much more extensive CI than Mohan et al. Gradually increasing the amount of CI, we have performed a series of calculations with the GRASP code, but focus only on our final results which include CI among 38 configurations (GRASP2), listed in Table 1. These configurations involve the $n \leq 5$ orbitals and generate 3749 levels in total. Their specific energy ranges are listed in Table 1, and the configurations included by Mohan et al. are also marked for reference. Two of the configurations, included by Mohan et al., namely $(3s^23p^3)\ 4d^2$ and $4f^2$, are not significant, because their levels lie in a very high energy range, i.e. $\geq$ 413 Ryd or $\sim$ 340 Ryd *above* the levels considered by these authors. Hence, their inclusion has little effect on the lower levels. On the other hand, Mohan et al. excluded some important configurations, such as $3s3p^33d^3$, $3s^23p3d^4$, and $3s3p^23d^4$. These configurations together generate 1716 levels and therefore significantly increase the size of a calculation, especially given that the 15 configurations of Mohan et al. generate only 1163 levels. However, the energy range of these three configurations is 92–196 Ryd, i.e. well below that of $3s^23p^44\ell$ included by

Mohan et al. There are additional configurations listed in Table 1 whose levels closely interact, intermix, and influence energies of the lower ones.

In Table 2 we compare our calculated energies for 31 levels of the $3s^23p^5$, $3s3p^6$, and $3s^23p^43d$ configurations with those of Mohan et al. [8]. Laboratory measurements have been compiled by Kramida [21] and the NIST (National Institute of Standards and Technology) team [22], which are also included in the table. As stated earlier, we have performed a series of calculations with increasing CI and one of these include the same 15 configurations (GRASP1) as considered by Mohan et al. Our energies for the levels obtained are comparable with their results listed in Table 2, but here we also list the mixing coefficients from the GRASP1 calculations. We note that Mohan et al. have listed wavefunction compositions (square of the mixing coefficient) in their Table 1, and hence provide a ready comparison. As a result of a larger CI included in the present work (GRASP2), the energies for a majority of levels are lower than the GRASP1 data by up to 0.07 Ryd, and hence are (slightly) closer to those of NIST. However, there are still discrepancies with the NIST listings of up to 0.2 Ryd for some levels, such as 13 ($3s^23p^4(^3P)3d\ ^2D_{5/2}$).

The NIST listings [22] for many levels are not based on direct measurements (see levels with a $\star$ in Table 2), but have been derived from interpolation and extrapolation. Similarly, two of their levels, i.e. $3s^23p^5\ ^2P^o_{1/2}$ and $3s^23p^4(^1S)3d\ ^2D_{5/2}$ (10 and 11) have the same energy, whereas in the theoretical work the difference between these is up to 0.1 Ryd, depending on the calculation. Finally, the NIST energy for the $3s3p^6\ ^2S_{1/2}$ level (14) is anomalous in comparison to all calculations – also see Table 2 of [8] and the present Table 3. For this level the NIST energy is *higher* by over 1 Ryd and needs a reassessment. Therefore, due to paucity of measured energy levels, the discrepancy noted with the NIST listings should not be considered an indication of the accuracy of our results.

To assess the accuracy of our energy levels, we have also performed parallel calculations with the *Flexible Atomic Code* (FAC) of Gu [23], available from the website `http://sprg.ssl.berkeley.edu/~mfgu/fac/`. This is also a fully relativistic code which provides results for energy levels and radiative rates comparable to GRASP, as already shown for several other ions, see for example: Aggarwal et al. for Kr [19] and Xe [20] ions, and more recent work on W XL by Aggarwal and Keenan [14]. With the same amount of CI as included in GRASP1 and GRASP2, the energy levels from FAC are comparable as listed in Table 2. This has also been shown by Mohan et al. [8] in their Table 2, and hence those results are not included in the present Table 2. However, we have performed four additional calculations with FAC, with increasing CI. These are (i) FAC1, which includes CI among 4580 levels of the 3*7 and $3s^23p^44\ell$ configurations, (ii) FAC2, which includes a further 1231 levels of $3s^23p^45\ell$, $3s3p^54\ell$, $3s3p^55\ell$, $3p^64\ell$, $3p^65\ell$, and $3s^23p^33d4\ell$, (iii) FAC3, with a total of 9160 levels, the additional ones arising from $3s^23p^46\ell$, $3s3p^56\ell$, $3p^66\ell$, $3s^23p^33d5\ell$, and $3s^23p^33d6\ell$, and finally (iv) FAC4, with a further 9299 levels (18,459 in total) of $3s^23p^34\ell^2$, $3s^23p^35\ell^2$, $3s^23p^34\ell5\ell$, $3p^54\ell^2$, and $3p^53d4\ell$. We discuss these results below.

In Table 3, we compare our GRASP2 energies with the four calculations from FAC described above. Also listed are the NIST [22] compilation for a ready reference. All calculations provide (nearly) the same orderings, and the differences between GRASP2 and all FAC calculations are less than ~ 0.1 Ryd, indicating that the CI included in our GRASP2 calculations is sufficient as far as the levels of Table 3 are concerned. Generally, the energies obtained with FAC are lower (by up to ~ 0.1 Ryd), as has also been noted by [8]. Based on these and earlier comparisons among a variety of calculations with two independent atomic structure codes and with increasing amount of CI, we assess the accuracy of our energy levels to be ~ 0.1 Ryd, or equivalently 0.5%.

Finally, we note that some of the levels are highly mixed as shown in the present Table 2 and in Table 1 of Mohan et al. [8]. Therefore, it is not always possible to provide a unique label for each level, but care has been taken to identify the levels as accurately as possible. Particularly difficult to identify are those levels which are highly mixed, such as 2/5/7/11/12/13 shown in Table 2. Therefore, the best one can say about a level is that it has a particular J value, as listed in Tables 2 and 3, but there can be disagreements about the configuration assigned to it. For this reason we differ in the designation of a few levels with those of Mohan et al., and examples include levels 5 and 7, i.e. ($3s^23p^4(^3P)3d$) $^2F_{7/2}$ and $^4D_{7/2}$. Ralchenko et al. [4] have the same

designations for these levels, but Mohan et al. have identified these as $3s^2 3p^4(^3P)3d\ ^4F_{7/2}$ and $3s^2 3p^4(^1D)3d\ ^2F_{7/2}$. This is a common problem for many ions in which level mixing is very strong, such as Ti VI [15], Cr VIII [16], Fe X [17], and Co XI [18].

3 Radiative rates

The absorption oscillator strength (f_{ij}), a dimensionless quantity, and radiative rate A_{ji} (in s^{-1}) for a transition $i \rightarrow j$ are related by the following expression:

$$f_{ij} = \frac{mc}{8\pi^2 e^2}\lambda_{ji}^2 \frac{\omega_j}{\omega_i} A_{ji} = 1.49 \times 10^{-16} \lambda_{ji}^2 \frac{\omega_j}{\omega_i} A_{ji} \tag{1}$$

where m and e are the electron mass and charge, respectively, c the velocity of light, λ_{ji} the transition energy/wavelength in Å, and ω_i and ω_j the statistical weights of the lower i and upper j levels, respectively. This relationship between the A- and f- values is the same irrespective of the type of a transition, such as electric dipole (E1), electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2). In Table 4 we list our A- values for all four types of transitions among the 31 levels of the $3s^2 3p^5$, $3s3p^6$, and $3s^2 3p^4 3d$ configurations, listed in Table 2. In addition, we list the f- values and line strengths (S- values), but for E1 transitions alone. For other transitions these can be easily determined though Eqs. (1–5) of [15].

The A- and f- values have been calculated in both Babushkin and Coulomb gauges, which are equivalent to the length and velocity forms in the non-relativistic nomenclature. However, the data in Table 4 are presented in the length form alone, as these are considered to be comparatively more accurate. Furthermore, these results correspond to our GRASP2 calculations and the differences (if any) with the corresponding GRASP1 results reported by Mohan et al. [8] are not significant for most of the E1 transitions. Indeed the only transition for which the two sets of A- values differ substantially is 1–11 ($3s^2 3p^5\ ^2P^o_{3/2}$ – $3s^2 3p^4(^1S)3d\ ^2D_{5/2}$), for which A (GRASP2) = 1.65×10^8 s^{-1} and A (GRASP1) = 7.57×10^7 s^{-1}. However, this is a *weak* transition with f $\sim 10^{-5}$. More importantly, Mohan et al. have reported A- values for only 31 E1 and 40 M2 transitions whereas there are 658 possible transitions, among all four types, i.e. 36 E1, 334 E2, 245 M1, and 43 M2, for which results are listed in Table 4. Finally, there is no discrepancy for the A- values of the 1–2 and 1–3 E1, and 3–7, 5–7, and 5–8 M1 transitions reported by Ralchenko et al. [4]. Based on the comparisons between the GRASP1 and GRASP2 calculations, as well as with FAC (see also Table 5 of [8]), we estimate the accuracy of our A-values and other related parameters to be better than 20%. This accuracy estimate is further confirmed by the velocity and length ratios of A- values for E1 transitions – see Table 3 of [8].

4 Lifetimes

The lifetime τ of a level j is defined as follows:

$$\tau_j = \frac{1}{\sum_i A_{ji}}. \tag{2}$$

In Table 5 we list lifetimes (τ) for all 31 levels from our calculations with the GRASP code. Results corresponding to our GRASP1 and GRASP2 calculations are listed, and *include* A- values from all types of transitions. Also listed are the lifetimes of Mohan et al. [8], which correspond to the GRASP1 calculations. As in the present work, they have included the contribution of A- values from all four types of transitions. The discrepancies between our calculated lifetimes and those of Mohan et al. are up to four orders of magnitude for about 30% of the levels, such as 8, 18, 22, 25, and 30. Although our GRASP2 calculations include a larger CI, the τ from our GRASP1 are comparable for all levels, except one, namely $3s^2 3p^4(^1S)3d\ ^2D_{5/2}$ (11). This is a direct consequence of the corresponding difference in the A- values of the 1–11 E1 transition, noted in section 3.

To understand the differences in τ between our calculations and those of Mohan et al. [8], we also list in Table 5 the A- values for the *dominant* transitions. The only level for which the E1 transition dominates and the difference in τ between the two calculations is four orders of magnitude is 30 ($3s^2 3p^4(^3P)3d\ ^4F_{3/2}$). For this level, the A- value for our 10–30 E1 transition from GRASP2 is 2.48×10^{12} s^{-1}, the same as from GRASP1, although Mohan et al. have not listed an A- value for this transition. Therefore, their reported τ of 1.10×10^{-9} s is unrealistic. Similarly, for level 18 ($3s^2 3p^4(^1D)3d\ ^2G_{7/2}$), the 5–18 M1 transition dominates with A = 1.60×10^{8} s^{-1} and $\tau = 4.6\times10^{-9}$ s , compared to $\tau = 5.7\times10^{-5}$ s of Mohan et al. Therefore, based on this discussion and the comparisons shown in Table 5, we can confidently state that some of the lifetimes reported by Mohan et al. are overestimated. Finally, there are no measurements available with which to compare our results, but we hope our reported lifetimes will be helpful to experimentalists.

5 Conclusions

In this work, energy levels, radiative rates, oscillator strengths, and lifetimes have been calculated with the GRASP code for a large number of levels/transitions of W LVIII, but results have been presented only among the 31 levels of the $3s^2 3p^5$, $3s3p^6$, and $3s^2 3p^4 3d$ configurations, as similar data have recently been reported by Mohan et al. [8]. These authors included limited CI in the determination of atomic structure and, more importantly, reported A- values for E1 and M2 transitions only from the levels of the ground state. This limited data is not helpful for modelling of plasmas, as demonstrated in Fig. 2 and Table 6 of Del Zanna et al. [24] for a similar Cl-like ion Fe X. To be specific level populations deduced using a limited data may be different by up to a factor of five. Therefore, in our work A- values have been listed for all possible 658 E1, E2, M1, and M2 transitions, which should be useful for plasma modelling.

As a result of the increased CI included in our work, the accuracy of the energy levels has improved by up to 0.07 Ryd, and brings the theoretical results closer to measurements, although limited to a few levels alone. The accuracy of our data is assessed to be better than 0.5% (within 0.1 Ryd), based on a series of calculations performed not only with the GRASP code, but also with FAC. Similarly, the accuracy of our radiative rates is assessed to be better than 20% for a majority of transitions.

Although no significant discrepancies are observed for the A- values reported by Mohan et al. [8] for a limited number of transitions, their calculated lifetimes are in error for several levels, by up to four orders of magnitude. Finally, similar results for a larger number of levels and their corresponding transitions will be reported in a later paper.

Acknowledgment

KMA is thankful to AWE Aldermaston for financial support.

References

[1] K. Fournier. At. Data Nucl. Data Tables **68**, 1 (1998).

[2] P. Quinet. J. Phys. **B 44**, 195007 (2011).

[3] S.B. Utter, P. Beiersdorfer, and E. Träbert. Can. J. Phys. **80**, 1503 (2002).

[4] Yu. Ralchenko, I. N. Draganic, J. N. Tan, J. D. Gillaspy, J. M. Pomeroy, J. Reader, U. Feldman, and G. E. Holland. J. Phys. **B 41**, 021003 (2008).

[5] J. Clementson, P. Beiersdorfer, G.V. Brown, M.F. Gu, H. Lundberg, Y. Podpaly, and E. Träbert. Can. J. Phys. **89**, 571 (2011).

[6] D.A. Liedahl. 2000 *Atomic Data Needs for X-ray Astronomy* available at `http://heasarc.gsfc.nasa.gov/docs/heasarc/atomic/`

[7] K.M. Aggarwal and F.P. Keenan. Fusion Sci. Tech. **63**, 363 (2013).

[8] M. Mohan, S. Aggarwal, and N. Singh. Can. J. Phys. **92**, xxx (2014), 10.1139/cjp-2013-0348.

[9] I.P. Grant, B.J. McKenzie, P.H. Norrington, D.F. Mayers, and N.C. Pyper. Comput. Phys. Commun. **21**, 207 (1980).

[10] K.G. Dyall, I.P. Grant, C.T. Johnson, F.A. Parpia, and E.P. Plummer. Comput. Phys. Commun., **55**, 425 (1989).

[11] F.A. Parpia, C.F. Fischer, and I.P. Grant. 1996 Comput. Phys. Commun. **94**, 249 (1996).

[12] P. Jönsson, X. He, C.F. Fischer, and I.P. Grant. 2007 Comput. Phys. Commun. **177**, 597 (2007).

[13] P. Jönsson P, G. Gaigalas, J. Bieroń, C.F. Fischer, and I.P. Grant. Comput. Phys. Commun. **184**, 2197 (2013).

[14] K.M. Aggarwal and F.P. Keenan. Can. J. Phys. **92**, xxx (2014).

[15] K.M. Aggarwal and F.P. Keenan. Phys. Scr. **88**, 025302 (2013).

[16] K.M. Aggarwal and F.P. Keenan. Astron. Astrophys. **506** (2009) 1501.

[17] K.M. Aggarwal and F.P. Keenan. Astron. Astrophys. **427**, 763 (2004).

[18] K.M. Aggarwal, F.P. Keenan, and A.Z. Msezane. Astron. Astrophys. **473**, 995 (2007).

[19] K.M. Aggarwal, F.P. Keenan, and K.D. Lawson. At. Data Nucl. Data Tables **94** 323 (2008).

[20] K.M. Aggarwal, F.P. Keenan, and K.D. Lawson. At. Data Nucl. Data Tables **96** 123 (2010).

[21] A. Kramida. Can. J. Phys. **89**, 551 (2011).

[22] A. Kramida, Y. Ralchenko, J. Reader, and NIST ASD Team. 2013 NIST Atomic Spectra Database (version 5.1) Online at: http://physics.nist.gov/asd

[23] M.F. Gu. Can. J. Phys. **86** 675 (2008).

[24] G. Del Zanna, K.A. Berrington, and H.E. Mason. Astron. Astrophys. **422**, 731 (2004).

Table 1.Configurations/Levels of W LVIII and their energy ranges (Ryd).

Index	Configuration	No. of Levels	Energy Range	GRASP1	GRASP2
1	$3s^23p^5$	2^o	0-26	Y	Y
2	$3s^23p^43d$	28	17-75	Y	Y
3	$3s3p^6$	1	40	Y	Y
4	$3s^23p^33d^2$	141^o	35-103	Y	Y
5	$3s3p^53d$	23^o	55-91	Y	Y
6	$3s^23p^23d^3$	261	55-130	Y	Y
7	$3s3p^43d^2$	211	73-143	Y	Y
8	$3s3p^33d^3$	678^o	92-169		Y
9	$3p^63d$	2	100-106	Y	Y
10	$3s^23p3d^4$	180^o	102-155		Y
11	$3s3p^23d^4$	858	113-196		Y
12	$3p^53d^2$	45^o	116-157	Y	Y
13	$3s^23p^44s$	8	182-209	Y	Y
14	$3s^23p^44p$	21^o	188-251	Y	Y
15	$3s^23p^44d$	28	205-261	Y	Y
16	$3s^23p^44f$	30^o	214-268	Y	Y
17	$3s^23p^33d4s$	72^o	199-262		Y
18	$3s^23p^33d4p$	203	205-278		Y
19	$3s^23p^33d4d$	302^o	222-288		Y
20	$3s^23p^33d4f$	363	231-295		Y
21	$3s3p^54s$	7^o	221-250		Y
22	$3s3p^54p$	18	227-265		Y
23	$3s3p^54d$	23^o	244-275		Y
24	$3s3p^54f$	24	253-283		Y
25	$3s^23p^45s$	8	276-329		Y
26	$3s^23p^45p$	21^o	279-337		Y
27	$3s^23p^45d$	28	288-342		Y
28	$3s^23p^45f$	30^o	292-346		Y
29	$3s^23p^45g$	30	295-348		Y
30	$3p^64s$	1	267		Y
31	$3p^64p$	2^o	271-283		Y
32	$3p^64d$	2	290-293		Y
33	$3p^64f$	2^o	298-300		Y
34	$3s3p^55s$	7	316-345		Y
35	$3s3p^55p$	18	320-353		Y
36	$3s3p^55d$	23^o	328-358		Y
37	$3s3p^55f$	24	332-361		Y
38	$3s3p^55g$	24^o	334-363		Y
39	$3s^23p^34d^2$	141^o	413-472	Y	
40	$3s^23p^34f^2$	221^o	430-487	Y	

GRASP1: earlier calculations of Mohan et al. [8] from the GRASP code with 1163 levels

GRASP2: present calculations from the GRASP code with 3749 levels

Y: configuration included under a calculation

Table 2. Energies (Ryd) for some levels of W LVIII and their mixing coefficients (MC).

Index	Configuration	Level	NIST	GRASP1	GRASP2	MC
1	$3s^23p^5$	$^2P^o_{3/2}$	00.000	0.0000	0.0000	1(0.995)
2	$3s^23p^4(^3P)3d$	$^4D_{3/2}$	17.344	17.4534	17.4004	2(-0.595)+16(0.455)+12(0.442)
3	$3s^23p^4(^3P)3d$	$^4D_{5/2}$	17.794	17.9141	17.8572	3(0.614)+19(0.447)
4	$3s^23p^4(^3P)3d$	$^4P_{1/2}$	17.870*	17.9752	17.9196	4(-0.708)+29(0.461)
5	$3s^23p^4(^3P)3d$	$^2F_{7/2}$	18.051	18.1905	18.1269	5(0.566)+22(0.539)+18(0.534)
6	$3s^23p^4(^1S)3d$	$^2D_{3/2}$	19.460*	19.6176	19.5587	6(0.758)+30(0.436)
7	$3s^23p^4(^3P)3d$	$^4D_{7/2}$	23.280	23.3713	23.3138	7(0.652)+27(0.502)+5(0.471)
8	$3s^23p^4(^3P)3d$	$^4F_{9/2}$	23.439	23.5406	23.4800	8(0.824)+25(0.562)
9	$3s^23p^4(^3P)3d$	$^2P_{1/2}$	24.080*	24.0296	23.9747	9(-0.672)+13(0.472)
10	$3s^23p^5$	$^2P^o_{1/2}$	25.500*	25.5348	25.5359	10(0.993)
11	$3s^23p^4(^1S)3d$	$^2D_{5/2}$	25.500*	25.6854	25.6227	11(0.681)+31(0.465)+24(-0.403)
12	$3s^23p^4(^3P)3d$	$^4P_{3/2}$	26.130*	26.1767	26.1047	12(-0.468)+23(0.453)+28(-0.444)
13	$3s^23p^4(^3P)3d$	$^2D_{5/2}$	26.600*	26.4839	26.4171	13(0.518)+3(-0.411)+11(-0.406)
14	$3s3p^6$	$^2S_{1/2}$	41.200*	39.8947	39.8704	14(0.855)
15	$3s^23p^4(^3P)3d$	$^4D_{1/2}$	42.200*	42.1254	42.0831	15(-0.886)
16	$3s^23p^4(^1D)3d$	$^2D_{3/2}$	42.900*	42.9204	42.8705	30(-0.571)+2(0.566)+16(0.311)
17	$3s^23p^4(^3P)3d$	$^4F_{5/2}$	43.500*	43.5241	43.4668	17(0.704)+19(0.460)
18	$3s^23p^4(^1D)3d$	$^2G_{7/2}$	43.670*	43.6284	43.5691	18(-0.770)+5(0.413)
19	$3s^23p^4(^1D)3d$	$^2F_{5/2}$	45.780	45.9809	45.9158	13(0.579)+19(-0.530)
20	$3s^23p^4(^1D)3d$	$^2P_{3/2}$	45.990	46.1664	46.0980	20(0.519)+16(-0.495)+12(0.474)
21	$3s^23p^4(^1D)3d$	$^2S_{1/2}$	46.450	46.6876	46.6326	21(0.517)+29(-0.508)+4(-0.472)
22	$3s^23p^4(^3P)3d$	$^4F_{7/2}$		48.3574	48.3091	7(0.663)+22(0.620)
23	$3s^23p^4(^3P)3d$	$^2P_{3/2}$		49.0564	49.0046	23(0.536)+12(-0.448)+20(-0.446)
24	$3s^23p^4(^3P)3d$	$^2F_{5/2}$		49.3626	49.3074	24(0.657)+31(0.473)
25	$3s^23p^4(^1D)3d$	$^2G_{9/2}$		49.3420	49.2842	25(-0.822)+8(0.561)
26	$3s^23p^4(^1D)3d$	$^2D_{5/2}$		49.8777	49.8205	26(-0.653)+31(0.520)
27	$3s^23p^4(^1D)3d$	$^2F_{7/2}$		50.5070	50.4437	27(-0.747)+5(0.436)
28	$3s^23p^4(^3P)3d$	$^2D_{3/2}$		52.2559	52.1908	28(-0.666)+16(-0.439)
29	$3s^23p^4(^1D)3d$	$^2P_{1/2}$		52.9774	52.9102	9(-0.571)+29(-0.557)+13(-0.547)
30	$3s^23p^4(^3P)3d$	$^4F_{3/2}$		71.0520	70.9979	6(0.585)+30(-0.504)+28(-0.405)
31	$3s^23p^4(^3P)3d$	$^4P_{5/2}$		75.4505	75.4010	11(0.589)+24(0.419)+31(-0.362)

NIST: [22], levels with a $\star$ are determined by interpolation/extrapolation

GRASP1: earlier calculations of Mohan et al. [8] from the GRASP code with 1163 levels

GRASP2: present calculations from the GRASP code with 3749 levels

MC: Mixing coefficients corresponding to GRASP1 calculations

Table 3. Comparison of energies (Ryd) from the GRASP and FAC codes for some levels of W LVIII.

Index	Configuration	Level	NIST	GRASP2	FAC1	FAC2	FAC3	FAC4
1	$3s^2 3p^5$	$^2P^o_{3/2}$	00.000	0.0000	0.00000	0.00000	0.00000	0.00000
2	$3s^2 3p^4(^3P)3d$	$^4D_{3/2}$	17.344	17.4004	17.32500	17.30007	17.29148	17.31476
3	$3s^2 3p^4(^3P)3d$	$^4D_{5/2}$	17.794	17.8572	17.78425	17.75610	17.74686	17.76984
4	$3s^2 3p^4(^3P)3d$	$^4P_{1/2}$	17.870	17.9196	17.84417	17.81809	17.80805	17.83068
5	$3s^2 3p^4(^3P)3d$	$^2F_{7/2}$	18.051	18.1269	18.05758	18.02481	18.01469	18.03738
6	$3s^2 3p^4(^1S)3d$	$^2D_{3/2}$	19.460	19.5587	19.48303	19.46102	19.45262	19.47488
7	$3s^2 3p^4(^3P)3d$	$^4D_{7/2}$	23.280	23.3138	23.25604	23.22210	23.21249	23.23596
8	$3s^2 3p^4(^3P)3d$	$^4F_{9/2}$	23.439	23.4800	23.42512	23.38844	23.37934	23.40281
9	$3s^2 3p^4(^3P)3d$	$^2P_{1/2}$	24.080	23.9747	23.91035	23.88147	23.87052	23.89290
10	$3s^2 3p^5$	$^2P^o_{1/2}$	25.500	25.5359	25.56424	25.56100	25.56023	25.56029
11	$3s^2 3p^4(^1S)3d$	$^2D_{5/2}$	25.500	25.6227	25.56139	25.53224	25.52054	25.54198
12	$3s^2 3p^4(^3P)3d$	$^4P_{3/2}$	26.130	26.1047	26.04906	26.01088	25.99531	26.01665
13	$3s^2 3p^4(^3P)3d$	$^2D_{5/2}$	26.600	26.4171	26.35345	26.32222	26.31125	26.33340
14	$3s3p^6$	$^2S_{1/2}$	41.200	39.8704	39.80383	39.79163	39.79341	39.80298
15	$3s^2 3p^4(^3P)3d$	$^4D_{1/2}$	42.200	42.0831	42.04051	42.01204	42.00635	42.03020
16	$3s^2 3p^4(^1D)3d$	$^2D_{3/2}$	42.900	42.8705	42.83200	42.79886	42.79085	42.81449
17	$3s^2 3p^4(^3P)3d$	$^4F_{5/2}$	43.500	43.4668	43.43225	43.39470	43.38525	43.40843
18	$3s^2 3p^4(^1D)3d$	$^2G_{7/2}$	43.670	43.5691	43.53531	43.49602	43.48692	43.51048
19	$3s^2 3p^4(^1D)3d$	$^2F_{5/2}$	45.780	45.9158	45.88080	45.84312	45.83029	45.85189
20	$3s^2 3p^4(^1D)3d$	$^2P_{3/2}$	45.990	46.0980	46.06912	46.02731	46.01176	46.03323
21	$3s^2 3p^4(^1D)3d$	$^2S_{1/2}$	46.450	46.6326	46.57652	46.55101	46.54240	46.56093
22	$3s^2 3p^4(^3P)3d$	$^4F_{7/2}$		48.3091	48.28499	48.24739	48.23930	48.26340
23	$3s^2 3p^4(^3P)3d$	$^2P_{3/2}$		49.0046	48.97824	48.94040	48.92859	48.95174
24	$3s^2 3p^4(^3P)3d$	$^2F_{5/2}$		49.3074	49.28477	49.24413	49.23220	49.25484
25	$3s^2 3p^4(^1D)3d$	$^2G_{9/2}$		49.2842	49.26511	49.22123	49.21161	49.23535
26	$3s^2 3p^4(^1D)3d$	$^2D_{5/2}$		49.8205	49.79443	49.75444	49.74101	49.76337
27	$3s^2 3p^4(^1D)3d$	$^2F_{7/2}$		50.4437	50.42117	50.37700	50.36281	50.38456
28	$3s^2 3p^4(^3P)3d$	$^2D_{3/2}$		52.1908	52.16259	52.12437	52.11111	52.13271
29	$3s^2 3p^4(^1D)3d$	$^2P_{1/2}$		52.9102	42.83200	42.79886	42.79085	42.81449
30	$3s^2 3p^4(^3P)3d$	$^4F_{3/2}$		70.9979	70.99515	70.95580	70.94608	70.96864
31	$3s^2 3p^4(^3P)3d$	$^4P_{5/2}$		75.4010	75.41228	75.37291	75.36308	75.38599

NIST: [22]

GRASP2: present calculations from the GRASP code with 3749 levels

FAC1: present calculations from the FAC code with 4580 levels

FAC2: present calculations from the FAC code with 5821 levels

FAC3: present calculations from the FAC code with 9160 levels

FAC4: present calculations from the FAC code with 18,459 levels

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
1	2	5.237+01	2.646+09	1.088−03	7.503−04	0.000+00	0.000+00	3.430+00
1	3	5.103+01	4.156+09	2.434−03	1.636−03	0.000+00	0.000+00	1.466+01
1	4	5.085+01	2.011+10	3.899−03	2.611−03	0.000+00	0.000+00	3.639+01
1	5	5.027+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	3.598+01
1	6	4.659+01	1.836+09	5.975−04	3.666−04	0.000+00	0.000+00	1.471+02
1	7	3.909+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	3.608+04
1	9	3.801+01	8.374+10	9.069−03	4.539−03	0.000+00	0.000+00	9.836+04
1	10	3.569+01	0.000+00	0.000+00	0.000+00	1.060+07	3.825+08	0.000+00
1	11	3.556+01	1.652+08	4.698−05	2.200−05	0.000+00	0.000+00	2.548+04
1	12	3.491+01	5.100+11	9.317−02	4.283−02	0.000+00	0.000+00	3.216+04
1	13	3.449+01	7.041+11	1.884−01	8.559−02	0.000+00	0.000+00	2.323+03
1	14	2.286+01	1.247+12	4.883−02	1.469−02	0.000+00	0.000+00	5.762+05
1	15	2.165+01	7.114+10	2.501−03	7.130−04	0.000+00	0.000+00	1.126+05
1	16	2.126+01	3.219+08	2.181−05	6.104−06	0.000+00	0.000+00	1.158+04
1	17	2.097+01	1.115+11	1.102−02	3.042−03	0.000+00	0.000+00	1.676+04
1	18	2.092+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	1.740+04
1	19	1.985+01	3.200+12	2.834−01	7.408−02	0.000+00	0.000+00	7.664+00
1	20	1.977+01	3.218+12	1.885−01	4.907−02	0.000+00	0.000+00	1.342+04
1	21	1.954+01	6.839+12	1.957−01	5.037−02	0.000+00	0.000+00	2.099+05
1	22	1.886+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	2.173+05
1	23	1.860+01	1.009+10	5.230−04	1.281−04	0.000+00	0.000+00	2.296+05
1	24	1.848+01	4.045+07	3.107−06	7.562−07	0.000+00	0.000+00	5.126+03
1	26	1.829+01	2.860+09	2.152−04	5.183−05	0.000+00	0.000+00	2.832+05
1	27	1.807+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	2.662+04
1	28	1.746+01	4.351+10	1.989−03	4.572−04	0.000+00	0.000+00	8.998+04
1	29	1.722+01	4.516+10	1.004−03	2.277−04	0.000+00	0.000+00	4.988+04
1	30	1.284+01	9.122+08	2.253−05	3.808−06	0.000+00	0.000+00	1.564+03
1	31	1.209+01	1.050+09	3.450−05	5.491−06	0.000+00	0.000+00	5.258+02
2	3	1.995+03	0.000+00	0.000+00	0.000+00	8.097−07	1.682+03	0.000+00
2	4	1.755+03	0.000+00	0.000+00	0.000+00	9.782−06	4.116+03	0.000+00
2	5	1.254+03	0.000+00	0.000+00	0.000+00	1.459−04	0.000+00	0.000+00
2	6	4.222+02	0.000+00	0.000+00	0.000+00	4.407+00	2.613+04	0.000+00
2	7	1.541+02	0.000+00	0.000+00	0.000+00	2.026+02	0.000+00	0.000+00
2	9	1.386+02	0.000+00	0.000+00	0.000+00	1.063+03	4.690+05	0.000+00
2	10	1.120+02	1.657+07	1.559−05	2.299−05	0.000+00	0.000+00	2.622−01
2	11	1.108+02	0.000+00	0.000+00	0.000+00	2.421+02	9.820+04	0.000+00
2	12	1.047+02	0.000+00	0.000+00	0.000+00	2.010+03	4.829+06	0.000+00
2	13	1.011+02	0.000+00	0.000+00	0.000+00	1.595−01	4.264+06	0.000+00
2	14	4.056+01	0.000+00	0.000+00	0.000+00	1.026+06	2.462+07	0.000+00
2	15	3.692+01	0.000+00	0.000+00	0.000+00	2.299+06	2.595+08	0.000+00

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
2	16	3.578+01	0.000+00	0.000+00	0.000+00	1.412+06	2.033+06	0.000+00
2	17	3.496+01	0.000+00	0.000+00	0.000+00	2.042+06	8.734+04	0.000+00
2	18	3.482+01	0.000+00	0.000+00	0.000+00	1.296+06	0.000+00	0.000+00
2	19	3.196+01	0.000+00	0.000+00	0.000+00	2.584+05	7.214+07	0.000+00
2	20	3.175+01	0.000+00	0.000+00	0.000+00	3.468+04	1.157+08	0.000+00
2	21	3.117+01	0.000+00	0.000+00	0.000+00	6.462+06	2.183+07	0.000+00
2	22	2.948+01	0.000+00	0.000+00	0.000+00	1.306+03	0.000+00	0.000+00
2	23	2.883+01	0.000+00	0.000+00	0.000+00	3.774+03	9.678+05	0.000+00
2	24	2.856+01	0.000+00	0.000+00	0.000+00	5.024+04	8.778+01	0.000+00
2	26	2.811+01	0.000+00	0.000+00	0.000+00	1.490+04	7.087+05	0.000+00
2	27	2.758+01	0.000+00	0.000+00	0.000+00	5.851+04	0.000+00	0.000+00
2	28	2.619+01	0.000+00	0.000+00	0.000+00	4.421+04	8.672+05	0.000+00
2	29	2.566+01	0.000+00	0.000+00	0.000+00	1.451+04	3.321+05	0.000+00
2	30	1.700+01	0.000+00	0.000+00	0.000+00	6.510+03	1.758+03	0.000+00
2	31	1.571+01	0.000+00	0.000+00	0.000+00	6.010+04	1.845+03	0.000+00
3	4	1.460+04	0.000+00	0.000+00	0.000+00	2.869−07	0.000+00	0.000+00
3	5	3.378+03	0.000+00	0.000+00	0.000+00	1.352−04	1.836+02	0.000+00
3	6	5.355+02	0.000+00	0.000+00	0.000+00	8.637+00	5.633+03	0.000+00
3	7	1.670+02	0.000+00	0.000+00	0.000+00	1.976+01	1.830+06	0.000+00
3	8	1.621+02	0.000+00	0.000+00	0.000+00	8.124+01	0.000+00	0.000+00
3	9	1.490+02	0.000+00	0.000+00	0.000+00	2.022+00	0.000+00	0.000+00
3	10	1.187+02	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	6.168−01
3	11	1.173+02	0.000+00	0.000+00	0.000+00	1.610+02	7.332+05	0.000+00
3	12	1.105+02	0.000+00	0.000+00	0.000+00	2.357+03	4.469+05	0.000+00
3	13	1.065+02	0.000+00	0.000+00	0.000+00	2.438+03	4.046+06	0.000+00
3	14	4.140+01	0.000+00	0.000+00	0.000+00	2.727+05	0.000+00	0.000+00
3	15	3.762+01	0.000+00	0.000+00	0.000+00	2.154+06	0.000+00	0.000+00
3	16	3.643+01	0.000+00	0.000+00	0.000+00	5.675+05	2.590+08	0.000+00
3	17	3.558+01	0.000+00	0.000+00	0.000+00	2.687+06	3.618+05	0.000+00
3	18	3.544+01	0.000+00	0.000+00	0.000+00	2.413+06	2.611+07	0.000+00
3	19	3.248+01	0.000+00	0.000+00	0.000+00	1.565+04	1.711+08	0.000+00
3	20	3.227+01	0.000+00	0.000+00	0.000+00	4.574+06	6.174+04	0.000+00
3	21	3.167+01	0.000+00	0.000+00	0.000+00	5.536+06	0.000+00	0.000+00
3	22	2.992+01	0.000+00	0.000+00	0.000+00	7.191+03	3.101+06	0.000+00
3	23	2.926+01	0.000+00	0.000+00	0.000+00	4.874+03	7.017+03	0.000+00
3	24	2.898+01	0.000+00	0.000+00	0.000+00	2.879+04	2.006+05	0.000+00
3	25	2.900+01	0.000+00	0.000+00	0.000+00	2.329+05	0.000+00	0.000+00
3	26	2.851+01	0.000+00	0.000+00	0.000+00	6.899+03	3.880+05	0.000+00
3	27	2.797+01	0.000+00	0.000+00	0.000+00	8.599+04	4.256+05	0.000+00
3	28	2.654+01	0.000+00	0.000+00	0.000+00	1.958+04	1.058+05	0.000+00

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
3	29	2.600+01	0.000+00	0.000+00	0.000+00	3.524+04	0.000+00	0.000+00
3	30	1.715+01	0.000+00	0.000+00	0.000+00	1.518+04	4.967+03	0.000+00
3	31	1.584+01	0.000+00	0.000+00	0.000+00	1.951+04	4.917+00	0.000+00
4	6	5.560+02	0.000+00	0.000+00	0.000+00	3.253+00	8.760+03	0.000+00
4	9	1.505+02	0.000+00	0.000+00	0.000+00	0.000+00	3.396+06	0.000+00
4	10	1.197+02	5.964+07	1.280−04	1.008−04	0.000+00	0.000+00	0.000+00
4	11	1.183+02	0.000+00	0.000+00	0.000+00	5.807+01	0.000+00	0.000+00
4	12	1.113+02	0.000+00	0.000+00	0.000+00	7.102+02	2.794+06	0.000+00
4	13	1.072+02	0.000+00	0.000+00	0.000+00	3.800+02	0.000+00	0.000+00
4	14	4.151+01	0.000+00	0.000+00	0.000+00	0.000+00	1.412+07	0.000+00
4	15	3.771+01	0.000+00	0.000+00	0.000+00	0.000+00	9.750+07	0.000+00
4	16	3.652+01	0.000+00	0.000+00	0.000+00	2.994+06	2.195+03	0.000+00
4	17	3.567+01	0.000+00	0.000+00	0.000+00	6.651+05	0.000+00	0.000+00
4	19	3.255+01	0.000+00	0.000+00	0.000+00	2.092+05	0.000+00	0.000+00
4	20	3.234+01	0.000+00	0.000+00	0.000+00	7.761+05	6.256+07	0.000+00
4	21	3.174+01	0.000+00	0.000+00	0.000+00	0.000+00	1.408+08	0.000+00
4	23	2.932+01	0.000+00	0.000+00	0.000+00	5.799+04	3.402+06	0.000+00
4	24	2.903+01	0.000+00	0.000+00	0.000+00	1.888+05	0.000+00	0.000+00
4	26	2.857+01	0.000+00	0.000+00	0.000+00	8.853+03	0.000+00	0.000+00
4	28	2.659+01	0.000+00	0.000+00	0.000+00	3.604+05	1.092+06	0.000+00
4	29	2.604+01	0.000+00	0.000+00	0.000+00	0.000+00	4.369+06	0.000+00
4	30	1.717+01	0.000+00	0.000+00	0.000+00	1.188+05	3.302+03	0.000+00
4	31	1.585+01	0.000+00	0.000+00	0.000+00	4.113+04	0.000+00	0.000+00
5	6	6.365+02	0.000+00	0.000+00	0.000+00	5.992+00	0.000+00	0.000+00
5	7	1.757+02	0.000+00	0.000+00	0.000+00	1.423+02	5.998+05	0.000+00
5	8	1.702+02	0.000+00	0.000+00	0.000+00	2.679+02	2.264+06	0.000+00
5	11	1.216+02	0.000+00	0.000+00	0.000+00	6.461+01	2.357+04	0.000+00
5	12	1.142+02	0.000+00	0.000+00	0.000+00	1.109+02	0.000+00	0.000+00
5	13	1.099+02	0.000+00	0.000+00	0.000+00	3.130+03	1.016+06	0.000+00
5	16	3.683+01	0.000+00	0.000+00	0.000+00	8.264+05	0.000+00	0.000+00
5	17	3.596+01	0.000+00	0.000+00	0.000+00	1.253+04	2.597+08	0.000+00
5	18	3.582+01	0.000+00	0.000+00	0.000+00	4.422+06	1.603+08	0.000+00
5	19	3.279+01	0.000+00	0.000+00	0.000+00	8.254+06	2.852+07	0.000+00
5	20	3.258+01	0.000+00	0.000+00	0.000+00	4.015+06	0.000+00	0.000+00
5	22	3.019+01	0.000+00	0.000+00	0.000+00	9.254+03	3.372+05	0.000+00
5	23	2.951+01	0.000+00	0.000+00	0.000+00	5.905+03	0.000+00	0.000+00
5	24	2.923+01	0.000+00	0.000+00	0.000+00	1.090+01	1.193+05	0.000+00
5	25	2.925+01	0.000+00	0.000+00	0.000+00	2.640+03	1.000+05	0.000+00
5	26	2.875+01	0.000+00	0.000+00	0.000+00	1.573+04	4.767+04	0.000+00
5	27	2.820+01	0.000+00	0.000+00	0.000+00	5.425+02	7.311+03	0.000+00

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
5	28	2.675+01	0.000+00	0.000+00	0.000+00	2.239+03	0.000+00	0.000+00
5	30	1.724+01	0.000+00	0.000+00	0.000+00	1.019+04	0.000+00	0.000+00
5	31	1.591+01	0.000+00	0.000+00	0.000+00	9.928+02	3.595+02	0.000+00
6	7	2.427+02	0.000+00	0.000+00	0.000+00	5.483−01	0.000+00	0.000+00
6	9	2.064+02	0.000+00	0.000+00	0.000+00	1.590+01	2.528+02	0.000+00
6	10	1.525+02	9.856+05	1.717−06	3.448−06	0.000+00	0.000+00	4.256−02
6	11	1.503+02	0.000+00	0.000+00	0.000+00	2.866+02	2.797+06	0.000+00
6	12	1.392+02	0.000+00	0.000+00	0.000+00	3.432+02	7.728+05	0.000+00
6	13	1.329+02	0.000+00	0.000+00	0.000+00	7.117+02	7.159+05	0.000+00
6	14	4.486+01	0.000+00	0.000+00	0.000+00	1.710+04	9.692+05	0.000+00
6	15	4.046+01	0.000+00	0.000+00	0.000+00	1.549+04	3.149+07	0.000+00
6	16	3.909+01	0.000+00	0.000+00	0.000+00	1.230+05	9.506+07	0.000+00
6	17	3.812+01	0.000+00	0.000+00	0.000+00	1.354+06	6.955+07	0.000+00
6	18	3.795+01	0.000+00	0.000+00	0.000+00	1.064+06	0.000+00	0.000+00
6	19	3.457+01	0.000+00	0.000+00	0.000+00	1.722+06	1.890+07	0.000+00
6	20	3.434+01	0.000+00	0.000+00	0.000+00	2.218+06	8.009+07	0.000+00
6	21	3.366+01	0.000+00	0.000+00	0.000+00	7.720+05	1.882+07	0.000+00
6	22	3.170+01	0.000+00	0.000+00	0.000+00	2.438+03	0.000+00	0.000+00
6	23	3.095+01	0.000+00	0.000+00	0.000+00	1.339+03	1.247+05	0.000+00
6	24	3.063+01	0.000+00	0.000+00	0.000+00	4.545+04	1.578+05	0.000+00
6	26	3.011+01	0.000+00	0.000+00	0.000+00	7.939+01	1.894+06	0.000+00
6	27	2.950+01	0.000+00	0.000+00	0.000+00	7.166+04	0.000+00	0.000+00
6	28	2.793+01	0.000+00	0.000+00	0.000+00	2.119+04	2.198+05	0.000+00
6	29	2.732+01	0.000+00	0.000+00	0.000+00	1.696+03	6.494+04	0.000+00
6	30	1.772+01	0.000+00	0.000+00	0.000+00	3.184+02	2.887+03	0.000+00
6	31	1.632+01	0.000+00	0.000+00	0.000+00	2.420+04	4.769+01	0.000+00
7	8	5.484+03	0.000+00	0.000+00	0.000+00	9.827−06	1.824+00	0.000+00
7	11	3.947+02	0.000+00	0.000+00	0.000+00	4.686+01	5.776+03	0.000+00
7	12	3.265+02	0.000+00	0.000+00	0.000+00	1.170+01	0.000+00	0.000+00
7	13	2.936+02	0.000+00	0.000+00	0.000+00	1.319+00	1.317+04	0.000+00
7	16	4.660+01	0.000+00	0.000+00	0.000+00	1.505+02	0.000+00	0.000+00
7	17	4.522+01	0.000+00	0.000+00	0.000+00	1.529+03	4.166+04	0.000+00
7	18	4.499+01	0.000+00	0.000+00	0.000+00	5.622+03	3.181+05	0.000+00
7	19	4.032+01	0.000+00	0.000+00	0.000+00	8.205+03	2.002+05	0.000+00
7	20	4.000+01	0.000+00	0.000+00	0.000+00	7.665+02	0.000+00	0.000+00
7	22	3.646+01	0.000+00	0.000+00	0.000+00	1.396+06	1.131+08	0.000+00
7	23	3.547+01	0.000+00	0.000+00	0.000+00	6.140+06	0.000+00	0.000+00
7	24	3.506+01	0.000+00	0.000+00	0.000+00	9.813+05	2.475+08	0.000+00
7	25	3.509+01	0.000+00	0.000+00	0.000+00	2.820+06	3.607+07	0.000+00
7	26	3.438+01	0.000+00	0.000+00	0.000+00	2.473+06	4.106+06	0.000+00

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
7	27	3.359+01	0.000+00	0.000+00	0.000+00	7.847+05	4.118+07	0.000+00
7	28	3.156+01	0.000+00	0.000+00	0.000+00	4.742+04	0.000+00	0.000+00
7	30	1.911+01	0.000+00	0.000+00	0.000+00	1.735+02	0.000+00	0.000+00
7	31	1.750+01	0.000+00	0.000+00	0.000+00	5.989+03	4.714+03	0.000+00
8	11	4.253+02	0.000+00	0.000+00	0.000+00	3.254+01	0.000+00	0.000+00
8	13	3.103+02	0.000+00	0.000+00	0.000+00	2.268+01	0.000+00	0.000+00
8	17	4.559+01	0.000+00	0.000+00	0.000+00	1.992+03	0.000+00	0.000+00
8	18	4.536+01	0.000+00	0.000+00	0.000+00	2.323+02	3.512+04	0.000+00
8	19	4.062+01	0.000+00	0.000+00	0.000+00	1.128+04	0.000+00	0.000+00
8	22	3.670+01	0.000+00	0.000+00	0.000+00	3.541+06	1.540+08	0.000+00
8	24	3.528+01	0.000+00	0.000+00	0.000+00	9.313+05	0.000+00	0.000+00
8	25	3.531+01	0.000+00	0.000+00	0.000+00	4.050+06	1.567+08	0.000+00
8	26	3.460+01	0.000+00	0.000+00	0.000+00	3.422+06	0.000+00	0.000+00
8	27	3.380+01	0.000+00	0.000+00	0.000+00	2.330+06	1.343+08	0.000+00
8	31	1.755+01	0.000+00	0.000+00	0.000+00	4.341+03	0.000+00	0.000+00
9	10	5.837+02	3.096+05	1.581−05	6.077−05	0.000+00	0.000+00	0.000+00
9	11	5.529+02	0.000+00	0.000+00	0.000+00	2.572+00	0.000+00	0.000+00
9	12	4.278+02	0.000+00	0.000+00	0.000+00	4.117+00	1.685+04	0.000+00
9	13	3.731+02	0.000+00	0.000+00	0.000+00	4.092−02	0.000+00	0.000+00
9	14	5.733+01	0.000+00	0.000+00	0.000+00	0.000+00	7.363+05	0.000+00
9	15	5.032+01	0.000+00	0.000+00	0.000+00	0.000+00	6.985+02	0.000+00
9	16	4.823+01	0.000+00	0.000+00	0.000+00	6.253+02	2.261+05	0.000+00
9	17	4.675+01	0.000+00	0.000+00	0.000+00	7.910+03	0.000+00	0.000+00
9	19	4.153+01	0.000+00	0.000+00	0.000+00	6.134+03	0.000+00	0.000+00
9	20	4.119+01	0.000+00	0.000+00	0.000+00	2.410+03	1.366+06	0.000+00
9	21	4.022+01	0.000+00	0.000+00	0.000+00	0.000+00	1.205+06	0.000+00
9	23	3.641+01	0.000+00	0.000+00	0.000+00	1.017+05	1.649+08	0.000+00
9	24	3.597+01	0.000+00	0.000+00	0.000+00	2.093+06	0.000+00	0.000+00
9	26	3.526+01	0.000+00	0.000+00	0.000+00	6.488+05	0.000+00	0.000+00
9	28	3.230+01	0.000+00	0.000+00	0.000+00	1.078+06	4.315+05	0.000+00
9	29	3.149+01	0.000+00	0.000+00	0.000+00	0.000+00	4.532+07	0.000+00
9	30	1.938+01	0.000+00	0.000+00	0.000+00	4.586+04	5.544+00	0.000+00
9	31	1.772+01	0.000+00	0.000+00	0.000+00	1.958+04	0.000+00	0.000+00
10	11	1.049+04	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	2.442−12
10	12	1.602+03	5.079+02	3.909−07	4.123−06	0.000+00	0.000+00	5.027−09
10	13	1.034+03	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	1.003−06
10	14	6.357+01	3.231+10	1.957−02	8.193−03	0.000+00	0.000+00	0.000+00
10	15	5.507+01	6.285+09	2.858−03	1.036−03	0.000+00	0.000+00	0.000+00
10	16	5.257+01	3.404+09	2.821−03	9.764−04	0.000+00	0.000+00	3.878+00
10	17	5.082+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	1.222+01

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
10	19	4.471+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	5.128+01
10	20	4.432+01	3.917+08	2.307−04	6.731−05	0.000+00	0.000+00	5.554+01
10	21	4.319+01	3.001+09	8.396−04	2.388−04	0.000+00	0.000+00	0.000+00
10	23	3.883+01	2.681+09	1.212−03	3.099−04	0.000+00	0.000+00	4.059+04
10	24	3.833+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	3.545+04
10	26	3.752+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	1.025+04
10	28	3.419+01	7.999+11	2.803−01	6.310−02	0.000+00	0.000+00	8.496+02
10	29	3.329+01	1.292+12	2.146−01	4.704−02	0.000+00	0.000+00	0.000+00
10	30	2.005+01	2.464+12	2.968−01	3.917−02	0.000+00	0.000+00	2.618+04
10	31	1.827+01	0.000+00	0.000+00	0.000+00	0.000+00	0.000+00	2.295+05
11	12	1.891+03	0.000+00	0.000+00	0.000+00	3.799−03	1.815+01	0.000+00
11	13	1.147+03	0.000+00	0.000+00	0.000+00	1.515−03	1.452+01	0.000+00
11	14	6.396+01	0.000+00	0.000+00	0.000+00	2.147+03	0.000+00	0.000+00
11	15	5.536+01	0.000+00	0.000+00	0.000+00	1.979+00	0.000+00	0.000+00
11	16	5.283+01	0.000+00	0.000+00	0.000+00	1.020+02	1.474+04	0.000+00
11	17	5.107+01	0.000+00	0.000+00	0.000+00	6.739+02	8.866+04	0.000+00
11	18	5.078+01	0.000+00	0.000+00	0.000+00	2.738+02	7.372+03	0.000+00
11	19	4.490+01	0.000+00	0.000+00	0.000+00	5.472+03	6.960+03	0.000+00
11	20	4.451+01	0.000+00	0.000+00	0.000+00	1.125+04	1.499+05	0.000+00
11	21	4.337+01	0.000+00	0.000+00	0.000+00	8.555+03	0.000+00	0.000+00
11	22	4.017+01	0.000+00	0.000+00	0.000+00	1.174+04	2.994+07	0.000+00
11	23	3.897+01	0.000+00	0.000+00	0.000+00	7.787+05	4.387+07	0.000+00
11	24	3.847+01	0.000+00	0.000+00	0.000+00	1.398+05	1.283+08	0.000+00
11	25	3.851+01	0.000+00	0.000+00	0.000+00	4.148+05	0.000+00	0.000+00
11	26	3.766+01	0.000+00	0.000+00	0.000+00	2.322+06	8.745+07	0.000+00
11	27	3.671+01	0.000+00	0.000+00	0.000+00	3.676+06	3.825+07	0.000+00
11	28	3.430+01	0.000+00	0.000+00	0.000+00	5.844+03	1.023+06	0.000+00
11	29	3.340+01	0.000+00	0.000+00	0.000+00	4.241+03	0.000+00	0.000+00
11	30	2.008+01	0.000+00	0.000+00	0.000+00	4.023+01	1.263+04	0.000+00
11	31	1.831+01	0.000+00	0.000+00	0.000+00	1.580+02	9.926+02	0.000+00
12	13	2.917+03	0.000+00	0.000+00	0.000+00	1.219−03	7.489+01	0.000+00
12	14	6.620+01	0.000+00	0.000+00	0.000+00	4.811+03	3.054+06	0.000+00
12	15	5.703+01	0.000+00	0.000+00	0.000+00	8.582+03	6.202+04	0.000+00
12	16	5.435+01	0.000+00	0.000+00	0.000+00	7.463+03	1.475+04	0.000+00
12	17	5.249+01	0.000+00	0.000+00	0.000+00	1.125+03	7.097+05	0.000+00
12	18	5.218+01	0.000+00	0.000+00	0.000+00	1.191+04	0.000+00	0.000+00
12	19	4.600+01	0.000+00	0.000+00	0.000+00	4.266+03	1.208+06	0.000+00
12	20	4.558+01	0.000+00	0.000+00	0.000+00	6.681+03	7.859+05	0.000+00
12	21	4.439+01	0.000+00	0.000+00	0.000+00	1.330+03	4.547+04	0.000+00
12	22	4.104+01	0.000+00	0.000+00	0.000+00	2.153+03	0.000+00	0.000+00

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
12	23	3.979+01	0.000+00	0.000+00	0.000+00	2.021+05	8.147+07	0.000+00
12	24	3.927+01	0.000+00	0.000+00	0.000+00	1.071+06	1.968+06	0.000+00
12	26	3.842+01	0.000+00	0.000+00	0.000+00	6.677+03	1.084+08	0.000+00
12	27	3.744+01	0.000+00	0.000+00	0.000+00	1.447+06	0.000+00	0.000+00
12	28	3.493+01	0.000+00	0.000+00	0.000+00	3.443+06	4.687+07	0.000+00
12	29	3.399+01	0.000+00	0.000+00	0.000+00	1.502+06	1.413+08	0.000+00
12	30	2.030+01	0.000+00	0.000+00	0.000+00	1.642+03	5.083+03	0.000+00
12	31	1.849+01	0.000+00	0.000+00	0.000+00	1.130+03	2.832+03	0.000+00
13	14	6.774+01	0.000+00	0.000+00	0.000+00	5.828+03	0.000+00	0.000+00
13	15	5.817+01	0.000+00	0.000+00	0.000+00	1.530+03	0.000+00	0.000+00
13	16	5.538+01	0.000+00	0.000+00	0.000+00	5.125+03	4.497+04	0.000+00
13	17	5.345+01	0.000+00	0.000+00	0.000+00	2.035+04	6.870+02	0.000+00
13	18	5.313+01	0.000+00	0.000+00	0.000+00	1.652+04	5.160+05	0.000+00
13	19	4.674+01	0.000+00	0.000+00	0.000+00	1.875−01	2.976+06	0.000+00
13	20	4.630+01	0.000+00	0.000+00	0.000+00	2.373+03	3.078+06	0.000+00
13	21	4.508+01	0.000+00	0.000+00	0.000+00	1.212+04	0.000+00	0.000+00
13	22	4.163+01	0.000+00	0.000+00	0.000+00	2.755+05	9.416+07	0.000+00
13	23	4.034+01	0.000+00	0.000+00	0.000+00	2.481+04	1.304+07	0.000+00
13	24	3.981+01	0.000+00	0.000+00	0.000+00	2.206+05	5.416+05	0.000+00
13	25	3.985+01	0.000+00	0.000+00	0.000+00	1.214+06	0.000+00	0.000+00
13	26	3.894+01	0.000+00	0.000+00	0.000+00	8.706+04	3.431+07	0.000+00
13	27	3.793+01	0.000+00	0.000+00	0.000+00	3.351+05	1.258+07	0.000+00
13	28	3.536+01	0.000+00	0.000+00	0.000+00	3.995+06	2.556+08	0.000+00
13	29	3.440+01	0.000+00	0.000+00	0.000+00	8.954+06	0.000+00	0.000+00
13	30	2.044+01	0.000+00	0.000+00	0.000+00	2.978+03	2.171+04	0.000+00
13	31	1.860+01	0.000+00	0.000+00	0.000+00	1.097+01	1.304+02	0.000+00
14	15	4.118+02	0.000+00	0.000+00	0.000+00	0.000+00	4.251+04	0.000+00
14	16	3.037+02	0.000+00	0.000+00	0.000+00	2.186+00	8.037+03	0.000+00
14	17	2.534+02	0.000+00	0.000+00	0.000+00	8.301−01	0.000+00	0.000+00
14	19	1.507+02	0.000+00	0.000+00	0.000+00	3.884−02	0.000+00	0.000+00
14	20	1.463+02	0.000+00	0.000+00	0.000+00	6.870+01	3.105+05	0.000+00
14	21	1.348+02	0.000+00	0.000+00	0.000+00	0.000+00	8.005+02	0.000+00
14	23	9.976+01	0.000+00	0.000+00	0.000+00	2.173+02	7.749+05	0.000+00
14	24	9.656+01	0.000+00	0.000+00	0.000+00	8.875+02	0.000+00	0.000+00
14	26	9.158+01	0.000+00	0.000+00	0.000+00	1.476+02	0.000+00	0.000+00
14	28	7.396+01	0.000+00	0.000+00	0.000+00	9.363+02	3.051+05	0.000+00
14	29	6.988+01	0.000+00	0.000+00	0.000+00	0.000+00	4.548+05	0.000+00
14	30	2.927+01	0.000+00	0.000+00	0.000+00	1.665+06	6.758+05	0.000+00
14	31	2.565+01	0.000+00	0.000+00	0.000+00	8.438+05	0.000+00	0.000+00
15	16	1.157+03	0.000+00	0.000+00	0.000+00	3.844−03	9.968+03	0.000+00

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
15	17	6.585+02	0.000+00	0.000+00	0.000+00	2.886−01	0.000+00	0.000+00
15	19	2.378+02	0.000+00	0.000+00	0.000+00	5.438+00	0.000+00	0.000+00
15	20	2.270+02	0.000+00	0.000+00	0.000+00	2.607+01	6.652+03	0.000+00
15	21	2.003+02	0.000+00	0.000+00	0.000+00	0.000+00	1.719+04	0.000+00
15	23	1.317+02	0.000+00	0.000+00	0.000+00	3.039+02	1.276+06	0.000+00
15	24	1.261+02	0.000+00	0.000+00	0.000+00	4.039+02	0.000+00	0.000+00
15	26	1.178+02	0.000+00	0.000+00	0.000+00	3.931−01	0.000+00	0.000+00
15	28	9.016+01	0.000+00	0.000+00	0.000+00	7.938+02	5.112+06	0.000+00
15	29	8.417+01	0.000+00	0.000+00	0.000+00	0.000+00	1.487+05	0.000+00
15	30	3.152+01	0.000+00	0.000+00	0.000+00	1.044+05	1.267+08	0.000+00
15	31	2.735+01	0.000+00	0.000+00	0.000+00	1.370+05	0.000+00	0.000+00
16	17	1.528+03	0.000+00	0.000+00	0.000+00	2.606−02	3.082+03	0.000+00
16	18	1.304+03	0.000+00	0.000+00	0.000+00	6.478−03	0.000+00	0.000+00
16	19	2.992+02	0.000+00	0.000+00	0.000+00	1.212−01	1.366+04	0.000+00
16	20	2.824+02	0.000+00	0.000+00	0.000+00	1.272+01	6.201+00	0.000+00
16	21	2.422+02	0.000+00	0.000+00	0.000+00	2.043+01	1.596+04	0.000+00
16	22	1.676+02	0.000+00	0.000+00	0.000+00	2.638+01	0.000+00	0.000+00
16	23	1.486+02	0.000+00	0.000+00	0.000+00	9.530+01	2.075+04	0.000+00
16	24	1.416+02	0.000+00	0.000+00	0.000+00	1.021+02	2.775+06	0.000+00
16	26	1.311+02	0.000+00	0.000+00	0.000+00	9.478+01	1.703+03	0.000+00
16	27	1.203+02	0.000+00	0.000+00	0.000+00	2.144+02	0.000+00	0.000+00
16	28	9.777+01	0.000+00	0.000+00	0.000+00	3.831+03	4.755+06	0.000+00
16	29	9.077+01	0.000+00	0.000+00	0.000+00	3.668+02	1.684+05	0.000+00
16	30	3.240+01	0.000+00	0.000+00	0.000+00	1.406+06	2.038+08	0.000+00
16	31	2.801+01	0.000+00	0.000+00	0.000+00	5.919+04	3.126+06	0.000+00
17	18	8.916+03	0.000+00	0.000+00	0.000+00	5.583−06	1.054+01	0.000+00
17	19	3.721+02	0.000+00	0.000+00	0.000+00	1.143+00	8.384+03	0.000+00
17	20	3.463+02	0.000+00	0.000+00	0.000+00	2.761+00	2.675+04	0.000+00
17	21	2.879+02	0.000+00	0.000+00	0.000+00	1.779+01	0.000+00	0.000+00
17	22	1.882+02	0.000+00	0.000+00	0.000+00	7.580+01	4.736+05	0.000+00
17	23	1.646+02	0.000+00	0.000+00	0.000+00	5.673−01	1.688+03	0.000+00
17	24	1.560+02	0.000+00	0.000+00	0.000+00	2.096+02	6.934+05	0.000+00
17	25	1.566+02	0.000+00	0.000+00	0.000+00	7.002+01	0.000+00	0.000+00
17	26	1.434+02	0.000+00	0.000+00	0.000+00	1.508+01	1.684+05	0.000+00
17	27	1.306+02	0.000+00	0.000+00	0.000+00	2.696+02	2.625+06	0.000+00
17	28	1.045+02	0.000+00	0.000+00	0.000+00	4.610+03	1.395+06	0.000+00
17	29	9.650+01	0.000+00	0.000+00	0.000+00	6.494+02	0.000+00	0.000+00
17	30	3.310+01	0.000+00	0.000+00	0.000+00	3.226+06	2.574+08	0.000+00
17	31	2.854+01	0.000+00	0.000+00	0.000+00	1.505+03	1.914+05	0.000+00
18	19	3.883+02	0.000+00	0.000+00	0.000+00	1.195+01	6.545+01	0.000+00

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
18	20	3.603+02	0.000+00	0.000+00	0.000+00	4.496+00	0.000+00	0.000+00
18	22	1.922+02	0.000+00	0.000+00	0.000+00	1.028+01	2.661+04	0.000+00
18	23	1.676+02	0.000+00	0.000+00	0.000+00	5.703+00	0.000+00	0.000+00
18	24	1.588+02	0.000+00	0.000+00	0.000+00	2.588+01	4.310+04	0.000+00
18	25	1.594+02	0.000+00	0.000+00	0.000+00	4.167+02	2.612+06	0.000+00
18	26	1.458+02	0.000+00	0.000+00	0.000+00	1.497+02	9.400+04	0.000+00
18	27	1.326+02	0.000+00	0.000+00	0.000+00	6.433+02	8.938+05	0.000+00
18	28	1.057+02	0.000+00	0.000+00	0.000+00	1.531+03	0.000+00	0.000+00
18	30	3.322+01	0.000+00	0.000+00	0.000+00	1.207+07	0.000+00	0.000+00
18	31	2.863+01	0.000+00	0.000+00	0.000+00	9.283+03	2.211+04	0.000+00
19	20	5.003+03	0.000+00	0.000+00	0.000+00	1.473−04	4.126+01	0.000+00
19	21	1.271+03	0.000+00	0.000+00	0.000+00	8.190−04	0.000+00	0.000+00
19	22	3.808+02	0.000+00	0.000+00	0.000+00	2.140+00	1.075+05	0.000+00
19	23	2.950+02	0.000+00	0.000+00	0.000+00	4.609+00	2.584+04	0.000+00
19	24	2.687+02	0.000+00	0.000+00	0.000+00	1.577−01	3.349+02	0.000+00
19	25	2.705+02	0.000+00	0.000+00	0.000+00	2.380+00	0.000+00	0.000+00
19	26	2.334+02	0.000+00	0.000+00	0.000+00	2.470+01	2.327+05	0.000+00
19	27	2.013+02	0.000+00	0.000+00	0.000+00	5.397+00	2.244+05	0.000+00
19	28	1.452+02	0.000+00	0.000+00	0.000+00	2.148+02	2.783+05	0.000+00
19	29	1.303+02	0.000+00	0.000+00	0.000+00	1.195+03	0.000+00	0.000+00
19	30	3.633+01	0.000+00	0.000+00	0.000+00	3.732+06	1.472+08	0.000+00
19	31	3.091+01	0.000+00	0.000+00	0.000+00	5.114+01	1.617+04	0.000+00
20	21	1.704+03	0.000+00	0.000+00	0.000+00	1.069−02	1.802+03	0.000+00
20	22	4.121+02	0.000+00	0.000+00	0.000+00	4.814−01	0.000+00	0.000+00
20	23	3.135+02	0.000+00	0.000+00	0.000+00	2.081+00	1.324+05	0.000+00
20	24	2.839+02	0.000+00	0.000+00	0.000+00	1.168+00	5.730+03	0.000+00
20	26	2.448+02	0.000+00	0.000+00	0.000+00	1.304+00	3.484+05	0.000+00
20	27	2.097+02	0.000+00	0.000+00	0.000+00	5.708+00	0.000+00	0.000+00
20	28	1.496+02	0.000+00	0.000+00	0.000+00	6.014+01	1.170+05	0.000+00
20	29	1.338+02	0.000+00	0.000+00	0.000+00	1.354+03	1.700+06	0.000+00
20	30	3.660+01	0.000+00	0.000+00	0.000+00	2.710+06	6.638+07	0.000+00
20	31	3.110+01	0.000+00	0.000+00	0.000+00	6.796+04	6.082+05	0.000+00
21	23	3.842+02	0.000+00	0.000+00	0.000+00	6.824−01	3.162+04	0.000+00
21	24	3.407+02	0.000+00	0.000+00	0.000+00	6.815−03	0.000+00	0.000+00
21	26	2.859+02	0.000+00	0.000+00	0.000+00	3.799+00	0.000+00	0.000+00
21	28	1.639+02	0.000+00	0.000+00	0.000+00	1.046+02	3.695+04	0.000+00
21	29	1.452+02	0.000+00	0.000+00	0.000+00	0.000+00	2.215+06	0.000+00
21	30	3.740+01	0.000+00	0.000+00	0.000+00	1.416+06	7.878+06	0.000+00
21	31	3.168+01	0.000+00	0.000+00	0.000+00	1.409+04	0.000+00	0.000+00
22	23	1.310+03	0.000+00	0.000+00	0.000+00	2.264−02	0.000+00	0.000+00

Table 4: Transition wavelengths in vacuum (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij}, dimensionless), and line strengths (S, in atomic units), in Babushkin gauge, for electric dipole (E1), and A_{ji} for E2, M1, and M2 transitions in W LVIII. ($a\pm b \equiv a\times 10^{\pm b}$).

i	j	λ_{ij}	A_{ji}^{E1}	f_{ij}^{E1}	S^{E1}	A_{ji}^{E2}	A_{ji}^{M1}	A_{ji}^{M2}
22	24	9.128+02	0.000+00	0.000+00	0.000+00	1.272−01	3.567+02	0.000+00
22	25	9.345+02	0.000+00	0.000+00	0.000+00	8.593−04	5.308+03	0.000+00
22	26	6.029+02	0.000+00	0.000+00	0.000+00	4.779+00	2.766+03	0.000+00
22	27	4.269+02	0.000+00	0.000+00	0.000+00	3.648+00	1.047+03	0.000+00
22	28	2.348+02	0.000+00	0.000+00	0.000+00	7.962−06	0.000+00	0.000+00
22	30	4.016+01	0.000+00	0.000+00	0.000+00	9.913+02	0.000+00	0.000+00
22	31	3.364+01	0.000+00	0.000+00	0.000+00	8.414+05	3.364+08	0.000+00
23	24	3.010+03	0.000+00	0.000+00	0.000+00	8.213−07	1.422+02	0.000+00
23	26	1.117+03	0.000+00	0.000+00	0.000+00	1.306−01	5.161+01	0.000+00
23	27	6.332+02	0.000+00	0.000+00	0.000+00	2.085−01	0.000+00	0.000+00
23	28	2.860+02	0.000+00	0.000+00	0.000+00	1.694−01	8.078+03	0.000+00
23	29	2.333+02	0.000+00	0.000+00	0.000+00	2.421+00	5.390+04	0.000+00
23	30	4.143+01	0.000+00	0.000+00	0.000+00	1.751+03	6.850+05	0.000+00
23	31	3.452+01	0.000+00	0.000+00	0.000+00	1.490+06	9.568+07	0.000+00
24	25	3.936+04	0.000+00	0.000+00	0.000+00	1.089−09	0.000+00	0.000+00
24	26	1.776+03	0.000+00	0.000+00	0.000+00	3.603−04	2.693+02	0.000+00
24	27	8.020+02	0.000+00	0.000+00	0.000+00	3.400−01	3.725+03	0.000+00
24	28	3.160+02	0.000+00	0.000+00	0.000+00	3.173−02	3.721+03	0.000+00
24	29	2.529+02	0.000+00	0.000+00	0.000+00	3.466+01	0.000+00	0.000+00
24	30	4.201+01	0.000+00	0.000+00	0.000+00	7.838+03	6.035+04	0.000+00
24	31	3.492+01	0.000+00	0.000+00	0.000+00	1.073+06	2.097+08	0.000+00
25	26	1.699+03	0.000+00	0.000+00	0.000+00	1.454−04	0.000+00	0.000+00
25	27	7.859+02	0.000+00	0.000+00	0.000+00	1.641+00	3.307+02	0.000+00
25	31	3.489+01	0.000+00	0.000+00	0.000+00	8.268+06	0.000+00	0.000+00
26	27	1.462+03	0.000+00	0.000+00	0.000+00	2.098−02	6.796+02	0.000+00
26	28	3.845+02	0.000+00	0.000+00	0.000+00	5.372+00	2.097+02	0.000+00
26	29	2.949+02	0.000+00	0.000+00	0.000+00	4.177+01	0.000+00	0.000+00
26	30	4.303+01	0.000+00	0.000+00	0.000+00	3.492+03	1.268+05	0.000+00
26	31	3.562+01	0.000+00	0.000+00	0.000+00	3.324+06	5.603+07	0.000+00
27	28	5.216+02	0.000+00	0.000+00	0.000+00	2.722+00	0.000+00	0.000+00
27	30	4.433+01	0.000+00	0.000+00	0.000+00	1.101+04	0.000+00	0.000+00
27	31	3.651+01	0.000+00	0.000+00	0.000+00	4.942+06	3.332+07	0.000+00
28	29	1.267+03	0.000+00	0.000+00	0.000+00	1.825−02	7.463+02	0.000+00
28	30	4.845+01	0.000+00	0.000+00	0.000+00	5.460+04	8.066+05	0.000+00
28	31	3.926+01	0.000+00	0.000+00	0.000+00	1.066+06	8.000+07	0.000+00
29	30	5.038+01	0.000+00	0.000+00	0.000+00	1.344+04	1.588+06	0.000+00
29	31	4.052+01	0.000+00	0.000+00	0.000+00	6.126+05	0.000+00	0.000+00
30	31	2.070+02	0.000+00	0.000+00	0.000+00	1.212+02	9.671+05	0.000+00

Table 5. Lifetimes (τ, s) for the levels of W LVIII and the *dominant* A- values. $a{\pm}b \equiv a{\times}10^{\pm b}$.

Index	Configuration	Level	GRASP1a	GRASP1b	GRASP2	Dominant A- values (s^{-1})
1	$3s^2 3p^5$	$^2P^o_{3/2}$	.....	.....	.....	.....
2	$3s^2 3p^4(^3P)3d$	$^4D_{3/2}$	3.73-10	3.737-10	3.779-10	1–2 E1=2.68+09
3	$3s^2 3p^4(^3P)3d$	$^4D_{5/2}$	2.40-10	2.399-10	2.406-10	1–3 E1=4.17+09
4	$3s^2 3p^4(^3P)3d$	$^4P_{1/2}$	4.89-11	4.898-11	4.972-11	1–4 E1=2.04+10
5	$3s^2 3p^4(^3P)3d$	$^2F_{7/2}$	2.65-02	4.269-03	4.555-03	3–5 E1=1.97+02, 1–5 M2=3.76+01
6	$3s^2 3p^4(^1S)3d$	$^2D_{3/2}$	5.49-10	5.498-10	5.447-10	1–6 E1=1.82+09
7	$3s^2 3p^4(^3P)3d$	$^4D_{7/2}$	2.74-05	4.053-07	4.054-07	3–7 M1=1.83+06, 1–7 M2=3.65+04
8	$3s^2 3p^4(^3P)3d$	$^4F_{9/2}$	5.59-03	4.418-07	4.415-07	5–8 M1=2.26+06, 5–8 E2=2.72+02
9	$3s^2 3p^4(^3P)3d$	$^2P_{1/2}$	1.18-11	1.181-11	1.194-11	1–9 E1=8.47+10
10	$3s^2 3p^5$	$^2P^o_{1/2}$	2.61-09	2.138-09	2.125-09	1–10 M1=3.83+08
11	$3s^2 3p^4(^1S)3d$	$^2D_{5/2}$	1.32-08	1.169-08	5.923-09	1–1 E1=8.19+07
12	$3s^2 3p^4(^3P)3d$	$^4P_{3/2}$	1.94-12	1.943-12	1.961-12	1–12 E1=5.15+11
13	$3s^2 3p^4(^3P)3d$	$^2D_{5/2}$	1.41-12	1.407-12	1.420-12	1–13 E1=7.11+11
14	$3s3p^6$	$^2S_{1/2}$	8.00-13	7.775-13	7.817-13	1–14 E1=1.25+12
15	$3s^2 3p^4(^3P)3d$	$^4D_{1/2}$	1.40-11	1.281-11	1.285-11	1–15 E1=7.14+10
16	$3s^2 3p^4(^1D)3d$	$^2D_{3/2}$	3.40-09	2.431-10	2.429-10	10–16 E1=3.42+09, 1–16 E1=2.97+08
17	$3s^2 3p^4(^3P)3d$	$^4F_{5/2}$	8.83-12	8.795-12	8.943-12	1–17 1.13+11
18	$3s^2 3p^4(^1D)3d$	$^2G_{7/2}$	5.71-05	4.617-09	4.656-09	5–18 M1=1.60+08, 3–18 M1=2.61+07
19	$3s^2 3p^4(^1D)3d$	$^2F_{5/2}$	3.10-13	3.105-13	3.125-13	1–19 E1=3.22+12
20	$3s^2 3p^4(^1D)3d$	$^2P_{3/2}$	3.08-13	3.085-13	3.107-13	1–20 E1=3.24+12
21	$3s^2 3p^4(^1D)3d$	$^2S_{1/2}$	1.46-13	1.455-13	1.462-13	1–21 E1=6.87+12
22	$3s^2 3p^4(^3P)3d$	$^4F_{7/2}$	4.57-06	2.431-09	2.435-09	7–22 M1=1.13+08, 8–22 M1=1.54+08
23	$3s^2 3p^4(^3P)3d$	$^2P_{3/2}$	9.56-11	7.414-11	7.629-11	1–23 E1=1.04+10, 10–23 E1=2.73+09
24	$3s^2 3p^4(^3P)3d$	$^2F_{5/2}$	2.21-08	2.223-09	2.252-09	7–24 M1=2.47+08, 11–24 M1=1.28+08
25	$3s^2 3p^4(^1D)3d$	$^2G_{9/2}$	4.32-06	4.050-09	4.123-09	8–25 M1=1.57+08, 7–25 M1=3.61+07
26	$3s^2 3p^4(^1D)3d$	$^2D_{5/2}$	3.42-10	3.124-10	3.168-10	1–26 E1=2.90+09
27	$3s^2 3p^4(^1D)3d$	$^2F_{7/2}$	3.73-05	3.369-09	3.432-09	8–27 M1=1.34+08, 7–27 M1=4.12+07
28	$3s^2 3p^4(^3P)3d$	$^2D_{3/2}$	2.26-11	1.172-12	1.185-12	10–28 E1=8.09+11
29	$3s^2 3p^4(^1D)3d$	$^2P_{1/2}$	2.34-11	7.414-13	7.479-13	10–29 E1=1.31+12
30	$3s^2 3p^4(^3P)3d$	$^4F_{3/2}$	1.10-09	4.022-13	4.054-13	10–30 E1=2.48+12
31	$3s^2 3p^4(^3P)3d$	$^4P_{5/2}$	2.83-10	2.495-10	2.574-10	1–31 E1=1.10+09

GRASP1a: earlier calculations of Mohan et al. [8] from the GRASP code with 1163 levels

GRASP1b: present calculations from the GRASP code with 1163 levels

GRASP2: present calculations from the GRASP code with 3749 levels